

## 6. The proximal mapping

- proximal mapping
- projections
- support functions, norms, distances

# Proximal mapping

**Definition:** the proximal mapping of a closed convex function  $f$  is

$$\text{prox}_f(x) = \underset{u}{\operatorname{argmin}} \left( f(u) + \frac{1}{2} \|u - x\|_2^2 \right)$$

**Existence and uniqueness:** we minimize a closed and strongly convex function

$$g(u) = f(u) + \frac{1}{2} \|u - x\|_2^2$$

- minimizer exists because  $g$  is closed with bounded sublevel sets
- minimizer is unique because  $g$  is strictly convex

**Subgradient characterization** (from page 4.7):

$$u = \text{prox}_f(x) \iff x - u \in \partial f(u)$$

# Examples

**Quadratic function** ( $A \geq 0$ )

$$f(x) = \frac{1}{2}x^T Ax + b^T x + c, \quad \text{prox}_{tf}(x) = (I + tA)^{-1}(x - tb)$$

**Euclidean norm:**  $f(x) = \|x\|_2$

$$\text{prox}_{tf}(x) = \begin{cases} (1 - t/\|x\|_2)x & \|x\|_2 \geq t \\ 0 & \text{otherwise} \end{cases}$$

**Logarithmic barrier**

$$f(x) = -\sum_{i=1}^n \log x_i, \quad \text{prox}_{tf}(x)_i = \frac{x_i + \sqrt{x_i^2 + 4t}}{2}, \quad i = 1, \dots, n$$

# Simple calculus rules

## Separable sum

$$f\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = g(x) + h(y), \quad \text{prox}_f\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} \text{prox}_g(x) \\ \text{prox}_h(y) \end{bmatrix}$$

**Scaling and translation of argument:** for scalar  $a \neq 0$ ,

$$f(x) = g(ax + b), \quad \text{prox}_f(x) = \frac{1}{a} \left( \text{prox}_{a^2 g}(ax + b) - b \right)$$

**“Right” scalar multiplication:** with  $\lambda > 0$ ,

$$f(x) = \lambda g(x/\lambda), \quad \text{prox}_f(x) = \lambda \text{prox}_{\lambda^{-1} g}(x/\lambda)$$

# Addition to linear or quadratic function

## Linear function

$$f(x) = g(x) + a^T x, \quad \text{prox}_f(x) = \text{prox}_g(x - a)$$

## Quadratic function: with $\mu > 0$

$$f(x) = g(x) + \frac{\mu}{2} \|x - a\|_2^2, \quad \text{prox}_f(x) = \text{prox}_{\theta g}(\theta x + (1 - \theta)a),$$

where  $\theta = 1/(1 + \mu)$

# Moreau decomposition

$$x = \text{prox}_f(x) + \text{prox}_{f^*}(x) \quad \text{for all } x$$

- follows from properties of conjugates and subgradients:

$$\begin{aligned} u = \text{prox}_f(x) &\Leftrightarrow x - u \in \partial f(u) \\ &\Leftrightarrow u \in \partial f^*(x - u) \\ &\Leftrightarrow x - u = \text{prox}_{f^*}(x) \end{aligned}$$

- generalizes decomposition by orthogonal projection on subspaces:

$$x = P_L(x) + P_{L^\perp}(x)$$

if  $L$  is a subspace,  $L^\perp$  its orthogonal complement

(this is the Moreau decomposition with  $f = \delta_L$ ,  $f^* = \delta_{L^\perp}$ )

## Extended Moreau decomposition

for  $\lambda > 0$ ,

$$x = \text{prox}_{\lambda f}(x) + \lambda \text{prox}_{\lambda^{-1} f^*}(x/\lambda) \quad \text{for all } x$$

*Proof:* apply Moreau decomposition to  $\lambda f$

$$\begin{aligned} x &= \text{prox}_{\lambda f}(x) + \text{prox}_{(\lambda f)^*}(x) \\ &= \text{prox}_{\lambda f}(x) + \lambda \text{prox}_{\lambda^{-1} f^*}(x/\lambda) \end{aligned}$$

second line uses  $(\lambda f)^*(y) = \lambda f^*(y/\lambda)$  and expression on page **6.4**

## Composition with affine mapping

$$f(x) = g(Ax + b)$$

- for general  $A$ , prox-operator of  $f$  does not follow easily from prox-operator of  $g$
- however, if  $AA^T = (1/\alpha)I$ , then

$$\begin{aligned}\text{prox}_f(x) &= (I - \alpha A^T A)x + \alpha A^T (\text{prox}_{\alpha^{-1}g}(Ax + b) - b) \\ &= x - \alpha A^T (Ax + b - \text{prox}_{\alpha^{-1}g}(Ax + b))\end{aligned}$$

**Example:**  $f(x_1, \dots, x_m) = g(x_1 + x_2 + \dots + x_m)$

- write as  $f(x) = g(Ax)$  with  $A = [ I \quad I \quad \dots \quad I ]$
- since  $AA^T = mI$ , we get

$$\text{prox}_f(x_1, \dots, x_m)_i = x_i - \frac{1}{m} \sum_{j=1}^m x_j + \frac{1}{m} \text{prox}_{mg}\left(\sum_{j=1}^m x_j\right), \quad i = 1, \dots, m$$



*Proof:*  $u = \text{prox}_f(x)$  is the solution of the optimization problem

$$\begin{aligned} & \text{minimize} && g(y) + \frac{1}{2}\|u - x\|_2^2 \\ & \text{subject to} && Au + b = y \end{aligned}$$

with variables  $u, y$

- eliminate  $u$  using the expression

$$\begin{aligned} u &= x + A^T(AA^T)^{-1}(y - b - Ax) \\ &= (I - \alpha A^T A)x + \alpha A^T(y - b) \quad (\text{since } AA^T = (1/\alpha)I) \end{aligned}$$

- optimal  $y$  is minimizer of

$$g(y) + \frac{\alpha^2}{2}\|A^T(y - b - Ax)\|_2^2 = g(y) + \frac{\alpha}{2}\|y - b - Ax\|_2^2$$

solution is  $y = \text{prox}_{\alpha^{-1}g}(Ax + b)$

# Outline

- conjugate functions
- proximal mapping
- **projections**
- support functions, norms, distances

# Projection on affine sets

**Hyperplane:**  $C = \{x \mid a^T x = b\}$  (with  $a \neq 0$ )

$$P_C(x) = x + \frac{b - a^T x}{\|a\|_2^2} a$$

**Affine set:**  $C = \{x \mid Ax = b\}$  (with  $A \in \mathbf{R}^{p \times n}$  and  $\mathbf{rank}(A) = p$ )

$$P_C(x) = x + A^T (AA^T)^{-1} (b - Ax)$$

inexpensive if  $p \ll n$ , or  $AA^T = I, \dots$

# Projection on simple polyhedral sets

**Halfspace:**  $C = \{x \mid a^T x \leq b\}$  (with  $a \neq 0$ )

$$P_C(x) = x + \frac{b - a^T x}{\|a\|_2^2} a \quad \text{if } a^T x > b, \quad P_C(x) = x \quad \text{if } a^T x \leq b$$

**Rectangle:**  $C = [l, u] = \{x \in \mathbf{R}^n \mid l \leq x \leq u\}$

$$P_C(x)_k = \begin{cases} l_k & x_k \leq l_k \\ x_k & l_k \leq x_k \leq u_k \\ u_k & x_k \geq u_k \end{cases}$$

**Nonnegative orthant:**  $C = \mathbf{R}_+^n$

$$P_C(x) = x_+ = (\max\{0, x_1\}, \max\{0, x_2\}, \dots, \max\{0, x_n\})$$

# Projection on simple polyhedral sets

**Probability simplex:**  $C = \{x \mid \mathbf{1}^T x = 1, x \geq 0\}$

$$P_C(x) = (x - \lambda \mathbf{1})_+$$

where  $\lambda$  is the solution of the equation

$$\mathbf{1}^T (x - \lambda \mathbf{1})_+ = \sum_{i=1}^n \max\{0, x_k - \lambda\} = 1$$

**Intersection of hyperplane and rectangle:**  $C = \{x \mid a^T x = b, l \leq x \leq u\}$

$$P_C(x) = P_{[l,u]}(x - \lambda a)$$

where  $\lambda$  is the solution of the equation

$$a^T P_{[l,u]}(x - \lambda a) = b$$

*Proof (probability simplex):* projection  $y = P_C(x)$  solves the optimization problem

$$\begin{aligned} & \text{minimize} && \frac{1}{2}\|y - x\|_2^2 + \delta_{\mathbf{R}_+^n}(y) \\ & \text{subject to} && \mathbf{1}^T y = 1 \end{aligned}$$

optimality conditions are:

- $y$  minimizes the Lagrangian

$$\begin{aligned} & \frac{1}{2}\|y - x\|_2^2 + \delta_{\mathbf{R}_+^n}(y) + \lambda(\mathbf{1}^T y - 1) \\ & = \sum_{k=1}^n \left( \frac{1}{2}(y_k - x_k)^2 + \delta_{\mathbf{R}_+}(y_k) + \lambda y_k \right) - \lambda \end{aligned}$$

this is a separable function with minimizer  $y_k = (x_k - \lambda)_+$  for  $k = 1, \dots, n$

- primal feasibility: requires

$$\sum_{k=1}^n y_i = \sum_{k=1}^n (x_k - \lambda)_+ = 1$$

*Proof (rectangle and hyperplane):*  $y = P_C(x)$  solves optimization problem

$$\begin{aligned} & \text{minimize} && \frac{1}{2}\|y - x\|_2^2 + \delta_{[l,u]}(y) \\ & \text{subject to} && a^T y = b \end{aligned}$$

optimality conditions are:

- $y$  minimizes the Lagrangian

$$\begin{aligned} & \frac{1}{2}\|y - x\|_2^2 + \delta_{[l,u]}(y) + \lambda(a^T y - b) \\ & = \sum_{k=1}^n \left( \frac{1}{2}(y_k - x_k)^2 + \delta_{[l_k, u_k]}(y_k) + \lambda a_k y_k \right) - \lambda b \end{aligned}$$

the minimizer is  $y_k = P_{[l_k, u_k]}(x_k - \lambda a_k)$  for  $k = 1, \dots, n$

- primal feasibility: requires

$$a^T y = \sum_{k=1}^n a_k P_{[l_k, u_k]}(x_k - \lambda a_k) = b$$

# Projection on norm balls

**Euclidean ball:**  $C = \{x \mid \|x\|_2 \leq 1\}$

$$P_C(x) = \frac{1}{\|x\|_2}x \quad \text{if } \|x\|_2 > 1, \quad P_C(x) = x \quad \text{if } \|x\|_2 \leq 1$$

**1-norm ball:**  $C = \{x \mid \|x\|_1 \leq 1\}$

projection is  $P_C(x) = x$  if  $\|x\|_1 \leq 1$ ; otherwise

$$P_C(x)_k = \text{sign}(x_k) \max\{|x_k| - \lambda, 0\} = \begin{cases} x_k - \lambda & x_k > \lambda \\ 0 & -\lambda \leq x_k \leq \lambda \\ x_k + \lambda & x_k < -\lambda \end{cases}$$

where  $\lambda$  is the solution of the equation

$$\sum_{k=1}^n \max\{|x_k| - \lambda, 0\} = 1$$



*Proof (1-norm):* projection  $y = P_C(x)$  solves the optimization problem

$$\begin{aligned} & \text{minimize} && \frac{1}{2}\|y - x\|_2^2 \\ & \text{subject to} && \|y\|_1 \leq 1 \end{aligned}$$

optimality conditions are:

- $y$  minimizes the Lagrangian

$$\frac{1}{2}\|y - x\|_2^2 + \lambda(\|y\|_1 - 1) = \sum_{k=1}^n \left( \frac{1}{2}(y_k - x_k)^2 + \lambda|y_k| \right) - \lambda$$

the minimizer  $y$  is obtained by componentwise soft-thresholding:

$$y_k = \text{sign}(x_k) \max\{|x_k| - \lambda, 0\}, \quad k = 1, \dots, n$$

- primal, dual feasibility and complementary slackness:

$$\lambda = 0, \quad \|y\|_1 = \|x\|_1 \leq 1 \quad \text{or} \quad \lambda > 0, \quad \|y\|_1 = \sum_{k=1}^n \max\{|x_k| - \lambda, 0\} = 1$$

## Projection on simple cones

**Second order cone:**  $C = \{(x, t) \in \mathbf{R}^{n \times 1} \mid \|x\|_2 \leq t\}$

$$P_C(x, t) = (x, t) \quad \text{if } \|x\|_2 \leq t, \quad P_C(x, t) = (0, 0) \quad \text{if } \|x\|_2 \leq -t$$

and

$$P_C(x, t) = \frac{t + \|x\|_2}{2\|x\|_2} \begin{bmatrix} x \\ \|x\|_2 \end{bmatrix} \quad \text{if } \|x\|_2 > |t|$$

**Positive semidefinite cone:**  $C = \mathbf{S}_+^n$

$$P_C(X) = \sum_{i=1}^n \max\{0, \lambda_i\} q_i q_i^T$$

if  $X = \sum_{i=1}^n \lambda_i q_i q_i^T$  is the eigenvalue decomposition of  $X$

# Outline

- conjugate functions
- proximal mapping
- projections
- **support functions, norms, distances**

# Support function

- conjugate of support function of closed convex set is indicator function

$$f(x) = \sup_{y \in C} x^T y, \quad f^*(y) = \delta_C(y)$$

- prox-operator of support function follows from Moreau decomposition

$$\begin{aligned} \text{prox}_t f(x) &= x - t \text{prox}_{t^{-1} f^*}(x/t) \\ &= x - t P_C(x/t) \end{aligned}$$

**Example:**  $f(x)$  is sum of largest  $r$  components of  $x$

$$f(x) = x_{[1]} + \cdots + x_{[r]} = \delta_C^*(x), \quad C = \{y \mid 0 \leq y \leq \mathbf{1}, \mathbf{1}^T y = r\}$$

prox-operator of  $f$  is easily evaluated via projection on  $C$  (page 6.12)

# Norms

- conjugate of norm is indicator function of dual norm ball:

$$f(x) = \|x\|, \quad f^*(y) = \delta_B(y) \quad \text{with } B = \{y \mid \|y\|_* \leq 1\}$$

- prox-operator of norm follows from Moreau decomposition

$$\begin{aligned} \text{prox}_t f(x) &= x - t \text{prox}_{t^{-1} f^*}(x/t) \\ &= x - t P_B(x/t) \\ &= x - P_{tB}(x) \end{aligned}$$

- gives  $\text{prox}_t \|\cdot\|$  when projection on  $tB = \{x \mid \|x\|_* \leq t\}$  is cheap

**Examples:** for  $\|\cdot\|_1$ ,  $\|\cdot\|_2$ , get expressions on pages 4.2 and 6.3

# Distance to a point

**Distance** (in general norm)

$$f(x) = \|x - a\|$$

**Prox-operator:** from page 6.4, with  $g(x) = \|x\|$

$$\begin{aligned}\text{prox}_{tf}(x) &= a + \text{prox}_{tg}(x - a) \\ &= a + x - a - tP_B\left(\frac{x - a}{t}\right) \\ &= x - P_{tB}(x - a)\end{aligned}$$

$B$  is the unit ball for the dual norm  $\|\cdot\|_*$

# Euclidean distance to a set

**Euclidean distance** (to a closed convex set  $C$ )

$$d(x) = \inf_{y \in C} \|x - y\|_2$$

**Prox-operator of distance**

$$\text{prox}_{td}(x) = \begin{cases} x + \frac{t}{d(x)}(P_C(x) - x) & d(x) \geq t \\ P_C(x) & \text{otherwise} \end{cases}$$


**Prox-operator of squared distance:**  $f(x) = d(x)^2/2$

$$\text{prox}_{tf}(x) = \frac{1}{1+t}x + \frac{t}{1+t}P_C(x)$$

*Proof* (expression for  $\text{prox}_{td}(x)$ ):

- if  $u = \text{prox}_{td}(x) \notin C$ , then from page 6.2 and subgradient for  $d$  (page 2.20)

$$x - u = \frac{t}{d(u)}(u - P_C(u))$$



- if  $\text{prox}_{td}(x) \in C$  then the minimizer of

$$d(u) + \frac{1}{2t} \|u - x\|_2^2$$

satisfies  $d(u) = 0$  and must be the projection  $P_C(x)$



*Proof* (expression for  $\text{prox}_{tf}(x)$  when  $f(x) = d(x)^2/2$ ):

$$\begin{aligned}\text{prox}_{tf}(x) &= \underset{u}{\text{argmin}} \left( \frac{1}{2}d(u)^2 + \frac{1}{2t}\|u - x\|_2^2 \right) \\ &= \underset{u}{\text{argmin}} \inf_{v \in C} \left( \frac{1}{2}\|u - v\|_2^2 + \frac{1}{2t}\|u - x\|_2^2 \right)\end{aligned}$$

- optimal  $u$  as a function of  $v$  is

$$u = \frac{t}{t+1}v + \frac{1}{t+1}x$$

- optimal  $v$  minimizes

$$\frac{1}{2} \left\| \frac{t}{t+1}v + \frac{1}{t+1}x - v \right\|_2^2 + \frac{1}{2t} \left\| \frac{t}{t+1}v + \frac{1}{t+1}x - x \right\|_2^2 = \frac{t}{2(1+t)} \|v - x\|_2^2$$

over  $C$ , *i.e.*,  $v = P_C(x)$

# References

- A. Beck, *First-Order Methods in Optimization* (2017), chapter 6.
- P. L. Combettes and J.-Ch. Pesquet, *Proximal splitting methods in signal processing*, in: *Fixed-Point Algorithms for Inverse Problems in Science and Engineering* (2011).
- N. Parikh and S. Boyd, *Proximal algorithms* (2013).