

5. Duality

- Lagrange dual problem
- weak and strong duality
- geometric interpretation
- optimality conditions
- perturbation and sensitivity analysis
- examples
- generalized inequalities

Lagrangian

standard form problem (not necessarily convex)

$$\begin{array}{ll} \text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & h_i(x) = 0, \quad i = 1, \dots, p \end{array}$$

variable $x \in \mathbf{R}^n$, domain \mathcal{D} , optimal value p^*

Lagrangian: $L : \mathbf{R}^n \times \mathbf{R}^m \times \mathbf{R}^p \rightarrow \mathbf{R}$, with $\text{dom } L = \mathcal{D} \times \mathbf{R}^m \times \mathbf{R}^p$,

$$L(x, \lambda, \nu) = f_0(x) + \sum_{i=1}^m \lambda_i f_i(x) + \sum_{i=1}^p \nu_i h_i(x)$$

- weighted sum of objective and constraint functions
- λ_i is Lagrange multiplier associated with $f_i(x) \leq 0$
- ν_i is Lagrange multiplier associated with $h_i(x) = 0$

Lagrange dual function

Lagrange dual function: $g : \mathbf{R}^m \times \mathbf{R}^p \rightarrow \mathbf{R}$,

$$\begin{aligned} g(\lambda, \nu) &= \inf_{x \in \mathcal{D}} L(x, \lambda, \nu) \\ &= \inf_{x \in \mathcal{D}} \left(f_0(x) + \sum_{i=1}^m \lambda_i f_i(x) + \sum_{i=1}^p \nu_i h_i(x) \right) \end{aligned}$$

g is concave, can be $-\infty$ for some λ, ν

lower bound property: if $\lambda \succeq 0$, then $g(\lambda, \nu) \leq p^*$

proof: if \tilde{x} is feasible and $\lambda \succeq 0$, then

$$f_0(\tilde{x}) \geq L(\tilde{x}, \lambda, \nu) \geq \inf_{x \in \mathcal{D}} L(x, \lambda, \nu) = g(\lambda, \nu)$$

minimizing over all feasible \tilde{x} gives $p^* \geq g(\lambda, \nu)$

Least-norm solution of linear equations

$$\begin{array}{ll} \text{minimize} & x^T x \\ \text{subject to} & Ax = b \end{array}$$

dual function

- Lagrangian is $L(x, \nu) = x^T x + \nu^T (Ax - b)$
- to minimize L over x , set gradient equal to zero:

$$\nabla_x L(x, \nu) = 2x + A^T \nu = 0 \quad \implies \quad x = -(1/2)A^T \nu$$

- plug in in L to obtain g :

$$g(\nu) = L((-1/2)A^T \nu, \nu) = -\frac{1}{4}\nu^T AA^T \nu - b^T \nu$$

a concave function of ν

lower bound property: $p^* \geq -(1/4)\nu^T AA^T \nu - b^T \nu$ for all ν

Standard form LP

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax = b, \quad x \succeq 0 \end{array}$$

dual function

- Lagrangian is

$$\begin{aligned} L(x, \lambda, \nu) &= c^T x + \nu^T (Ax - b) - \lambda^T x \\ &= -b^T \nu + (c + A^T \nu - \lambda)^T x \end{aligned}$$

- L is affine in x , hence

$$g(\lambda, \nu) = \inf_x L(x, \lambda, \nu) = \begin{cases} -b^T \nu & A^T \nu - \lambda + c = 0 \\ -\infty & \text{otherwise} \end{cases}$$

g is linear on affine domain $\{(\lambda, \nu) \mid A^T \nu - \lambda + c = 0\}$, hence concave

lower bound property: $p^* \geq -b^T \nu$ if $A^T \nu + c \succeq 0$

Equality constrained norm minimization

$$\begin{array}{ll} \text{minimize} & \|x\| \\ \text{subject to} & Ax = b \end{array}$$

dual function

$$g(\nu) = \inf_x (\|x\| - \nu^T Ax + b^T \nu) = \begin{cases} b^T \nu & \|A^T \nu\|_* \leq 1 \\ -\infty & \text{otherwise} \end{cases}$$

where $\|v\|_* = \sup_{\|u\| \leq 1} u^T v$ is dual norm of $\|\cdot\|$

proof: follows from $\inf_x (\|x\| - y^T x) = 0$ if $\|y\|_* \leq 1$, $-\infty$ otherwise

- if $\|y\|_* \leq 1$, then $\|x\| - y^T x \geq 0$ for all x , with equality if $x = 0$
- if $\|y\|_* > 1$, choose $x = tu$ where $\|u\| \leq 1$, $u^T y = \|y\|_* > 1$:

$$\|x\| - y^T x = t(\|u\| - \|y\|_*) \rightarrow -\infty \quad \text{as } t \rightarrow \infty$$

lower bound property: $p^* \geq b^T \nu$ if $\|A^T \nu\|_* \leq 1$

Two-way partitioning

$$\begin{array}{ll} \text{minimize} & x^T W x \\ \text{subject to} & x_i^2 = 1, \quad i = 1, \dots, n \end{array}$$

- a nonconvex problem; feasible set contains 2^n discrete points
- interpretation: partition $\{1, \dots, n\}$ in two sets; W_{ij} is cost of assigning i, j to the same set; $-W_{ij}$ is cost of assigning to different sets

dual function

$$\begin{aligned} g(\nu) &= \inf_x (x^T W x + \sum_i \nu_i (x_i^2 - 1)) = \inf_x x^T (W + \mathbf{diag}(\nu)) x - \mathbf{1}^T \nu \\ &= \begin{cases} -\mathbf{1}^T \nu & W + \mathbf{diag}(\nu) \succeq 0 \\ -\infty & \text{otherwise} \end{cases} \end{aligned}$$

lower bound property: $p^* \geq -\mathbf{1}^T \nu$ if $W + \mathbf{diag}(\nu) \succeq 0$

example: $\nu = -\lambda_{\min}(W)\mathbf{1}$ gives bound $p^* \geq n\lambda_{\min}(W)$

Lagrange dual and conjugate function

$$\begin{array}{ll} \text{minimize} & f_0(x) \\ \text{subject to} & Ax \preceq b, \quad Cx = d \end{array}$$

dual function

$$\begin{aligned} g(\lambda, \nu) &= \inf_{x \in \text{dom } f_0} (f_0(x) + (A^T \lambda + C^T \nu)^T x - b^T \lambda - d^T \nu) \\ &= -f_0^*(-A^T \lambda - C^T \nu) - b^T \lambda - d^T \nu \end{aligned}$$

- recall definition of conjugate $f^*(y) = \sup_{x \in \text{dom } f} (y^T x - f(x))$
- simplifies derivation of dual if conjugate of f_0 is known

example: entropy maximization

$$f_0(x) = \sum_{i=1}^n x_i \log x_i, \quad f_0^*(y) = \sum_{i=1}^n e^{y_i - 1}$$

The dual problem

Lagrange dual problem

$$\begin{array}{ll} \text{maximize} & g(\lambda, \nu) \\ \text{subject to} & \lambda \succeq 0 \end{array}$$

- finds best lower bound on p^* , obtained from Lagrange dual function
- a convex optimization problem; optimal value denoted d^*
- λ, ν are dual feasible if $\lambda \succeq 0, (\lambda, \nu) \in \mathbf{dom} g$
- often simplified by making implicit constraint $(\lambda, \nu) \in \mathbf{dom} g$ explicit

example: standard form LP and its dual (page 5–5)

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \succeq 0 \end{array}$$

$$\begin{array}{ll} \text{maximize} & -b^T \nu \\ \text{subject to} & A^T \nu + c \succeq 0 \end{array}$$

Weak and strong duality

weak duality: $d^* \leq p^*$

- always holds (for convex and nonconvex problems)
- can be used to find nontrivial lower bounds for difficult problems
for example, solving the SDP

$$\begin{array}{ll} \text{maximize} & -\mathbf{1}^T \nu \\ \text{subject to} & W + \mathbf{diag}(\nu) \succeq 0 \end{array}$$

gives a lower bound for the two-way partitioning problem on page 5–7

strong duality: $d^* = p^*$

- does not hold in general
- (usually) holds for convex problems
- conditions that guarantee strong duality in convex problems are called **constraint qualifications**

Slater's constraint qualification

strong duality holds for a convex problem

$$\begin{array}{ll} \text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & Ax = b \end{array}$$

if it is strictly feasible, *i.e.*,

$$\exists x \in \mathbf{int} \mathcal{D} : \quad f_i(x) < 0, \quad i = 1, \dots, m, \quad Ax = b$$

- also guarantees that the dual optimum is attained (if $p^* > -\infty$)
- can be sharpened: *e.g.*, can replace $\mathbf{int} \mathcal{D}$ with $\mathbf{relint} \mathcal{D}$ (interior relative to affine hull); linear inequalities do not need to hold with strict inequality, . . .
- there exist many other types of constraint qualifications

Inequality form LP

primal problem

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax \preceq b \end{array}$$

dual function

$$g(\lambda) = \inf_x ((c + A^T \lambda)^T x - b^T \lambda) = \begin{cases} -b^T \lambda & A^T \lambda + c = 0 \\ -\infty & \text{otherwise} \end{cases}$$

dual problem

$$\begin{array}{ll} \text{maximize} & -b^T \lambda \\ \text{subject to} & A^T \lambda + c = 0, \quad \lambda \succeq 0 \end{array}$$

- from Slater's condition: $p^* = d^*$ if $A\tilde{x} \prec b$ for some \tilde{x}
- in fact, $p^* = d^*$ except when primal and dual are infeasible

Quadratic program

primal problem (assume $P \in \mathbf{S}_{++}^n$)

$$\begin{aligned} & \text{minimize} && x^T P x \\ & \text{subject to} && Ax \preceq b \end{aligned}$$

dual function

$$g(\lambda) = \inf_x (x^T P x + \lambda^T (Ax - b)) = -\frac{1}{4} \lambda^T A P^{-1} A^T \lambda - b^T \lambda$$

dual problem

$$\begin{aligned} & \text{maximize} && -(1/4) \lambda^T A P^{-1} A^T \lambda - b^T \lambda \\ & \text{subject to} && \lambda \succeq 0 \end{aligned}$$

- from Slater's condition: $p^* = d^*$ if $A\tilde{x} \prec b$ for some \tilde{x}
- in fact, $p^* = d^*$ always

A nonconvex problem with strong duality

$$\begin{aligned} & \text{minimize} && x^T A x + 2b^T x \\ & \text{subject to} && x^T x \leq 1 \end{aligned}$$

$A \not\preceq 0$, hence nonconvex

dual function: $g(\lambda) = \inf_x (x^T (A + \lambda I)x + 2b^T x - \lambda)$

- unbounded below if $A + \lambda I \not\preceq 0$ or if $A + \lambda I \succeq 0$ and $b \notin \mathcal{R}(A + \lambda I)$
- minimized by $x = -(A + \lambda I)^\dagger b$ otherwise: $g(\lambda) = -b^T (A + \lambda I)^\dagger b - \lambda$

dual problem and equivalent SDP:

$$\begin{aligned} & \text{maximize} && -b^T (A + \lambda I)^\dagger b - \lambda \\ & \text{subject to} && A + \lambda I \succeq 0 \\ & && b \in \mathcal{R}(A + \lambda I) \end{aligned}$$

$$\begin{aligned} & \text{maximize} && -t - \lambda \\ & \text{subject to} && \begin{bmatrix} A + \lambda I & b \\ b^T & t \end{bmatrix} \succeq 0 \end{aligned}$$


strong duality although primal problem is not convex (not easy to show)

Geometric interpretation

for simplicity, consider problem with one constraint $f_1(x) \leq 0$

interpretation of dual function:


$$g(\lambda) = \inf_{(u,t) \in \mathcal{G}} (t + \lambda u), \quad \text{where } \mathcal{G} = \{(f_1(x), f_0(x)) \mid x \in \mathcal{D}\}$$



- $\lambda u + t = g(\lambda)$ is (non-vertical) supporting hyperplane to \mathcal{G}
- hyperplane intersects t -axis at $t = g(\lambda)$

epigraph variation: same interpretation if \mathcal{G} is replaced with

$$\mathcal{A} = \{(u, t) \mid f_1(x) \leq u, f_0(x) \leq t \text{ for some } x \in \mathcal{D}\}$$



strong duality

- holds if there is a non-vertical supporting hyperplane to \mathcal{A} at $(0, p^*)$
- for convex problem, \mathcal{A} is convex, hence has supp. hyperplane at $(0, p^*)$
- Slater's condition: if there exist $(\tilde{u}, \tilde{t}) \in \mathcal{A}$ with $\tilde{u} < 0$, then supporting hyperplanes at $(0, p^*)$ must be non-vertical

Complementary slackness

assume strong duality holds, x^* is primal optimal, (λ^*, ν^*) is dual optimal

$$\begin{aligned} f_0(x^*) = g(\lambda^*, \nu^*) &= \inf_x \left(f_0(x) + \sum_{i=1}^m \lambda_i^* f_i(x) + \sum_{i=1}^p \nu_i^* h_i(x) \right) \\ &\leq f_0(x^*) + \sum_{i=1}^m \lambda_i^* f_i(x^*) + \sum_{i=1}^p \nu_i^* h_i(x^*) \\ &\leq f_0(x^*) \end{aligned}$$

hence, the two inequalities hold with equality

- x^* minimizes $L(x, \lambda^*, \nu^*)$
- $\lambda_i^* f_i(x^*) = 0$ for $i = 1, \dots, m$ (known as complementary slackness):

$$\lambda_i^* > 0 \implies f_i(x^*) = 0, \quad f_i(x^*) < 0 \implies \lambda_i^* = 0$$

Karush-Kuhn-Tucker (KKT) conditions

the following four conditions are called KKT conditions (for a problem with differentiable f_i, h_i):

1. primal constraints: $f_i(x) \leq 0, i = 1, \dots, m, h_i(x) = 0, i = 1, \dots, p$
2. dual constraints: $\lambda \succeq 0$
3. complementary slackness: $\lambda_i f_i(x) = 0, i = 1, \dots, m$
4. gradient of Lagrangian with respect to x vanishes:

$$\nabla f_0(x) + \sum_{i=1}^m \lambda_i \nabla f_i(x) + \sum_{i=1}^p \nu_i \nabla h_i(x) = 0$$

from page 5–17: if strong duality holds and x, λ, ν are optimal, then they must satisfy the KKT conditions

KKT conditions for convex problem

if \tilde{x} , $\tilde{\lambda}$, $\tilde{\nu}$ satisfy KKT for a convex problem, then they are optimal:

- from complementary slackness: $f_0(\tilde{x}) = L(\tilde{x}, \tilde{\lambda}, \tilde{\nu})$
- from 4th condition (and convexity): $g(\tilde{\lambda}, \tilde{\nu}) = L(\tilde{x}, \tilde{\lambda}, \tilde{\nu})$

hence, $f_0(\tilde{x}) = g(\tilde{\lambda}, \tilde{\nu})$

if **Slater's condition** is satisfied:

x is optimal if and only if there exist λ , ν that satisfy KKT conditions

- recall that Slater implies strong duality, and dual optimum is attained
- generalizes optimality condition $\nabla f_0(x) = 0$ for unconstrained problem

example: water-filling (assume $\alpha_i > 0$)

$$\begin{aligned} & \text{minimize} && -\sum_{i=1}^n \log(x_i + \alpha_i) \\ & \text{subject to} && x \succeq 0, \quad \mathbf{1}^T x = 1 \end{aligned}$$


x is optimal iff $x \succeq 0$, $\mathbf{1}^T x = 1$, and there exist $\lambda \in \mathbf{R}^n$, $\nu \in \mathbf{R}$ such that

$$\lambda \succeq 0, \quad \lambda_i x_i = 0, \quad \frac{1}{x_i + \alpha_i} + \lambda_i = \nu$$

- if $\nu < 1/\alpha_i$: $\lambda_i = 0$ and $x_i = 1/\nu - \alpha_i$
- if $\nu \geq 1/\alpha_i$: $\lambda_i = \nu - 1/\alpha_i$ and $x_i = 0$
- determine ν from $\mathbf{1}^T x = \sum_{i=1}^n \max\{0, 1/\nu - \alpha_i\} = 1$

interpretation

- n patches; level of patch i is at height α_i
- flood area with unit amount of water
- resulting level is $1/\nu^*$



Perturbation and sensitivity analysis

(unperturbed) optimization problem and its dual

$$\begin{array}{ll} \text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & h_i(x) = 0, \quad i = 1, \dots, p \end{array} \qquad \begin{array}{ll} \text{maximize} & g(\lambda, \nu) \\ \text{subject to} & \lambda \succeq 0 \end{array}$$

perturbed problem and its dual

$$\begin{array}{ll} \text{min.} & f_0(x) \\ \text{s.t.} & f_i(x) \leq u_i, \quad i = 1, \dots, m \\ & h_i(x) = v_i, \quad i = 1, \dots, p \end{array} \qquad \begin{array}{ll} \text{max.} & g(\lambda, \nu) - u^T \lambda - v^T \nu \\ \text{s.t.} & \lambda \succeq 0 \end{array}$$

- x is primal variable; u, v are parameters
- $p^*(u, v)$ is optimal value as a function of u, v
- we are interested in information about $p^*(u, v)$ that we can obtain from the solution of the unperturbed problem and its dual

global sensitivity result

assume strong duality holds for unperturbed problem, and that λ^* , ν^* are dual optimal for unperturbed problem

apply weak duality to perturbed problem:

$$\begin{aligned} p^*(u, v) &\geq g(\lambda^*, \nu^*) - u^T \lambda^* - v^T \nu^* \\ &= p^*(0, 0) - u^T \lambda^* - v^T \nu^* \end{aligned}$$

sensitivity interpretation

- if λ_i^* large: p^* increases greatly if we tighten constraint i ($u_i < 0$)
- if λ_i^* small: p^* does not decrease much if we loosen constraint i ($u_i > 0$)
- if ν_i^* large and positive: p^* increases greatly if we take $v_i < 0$;
if ν_i^* large and negative: p^* increases greatly if we take $v_i > 0$
- if ν_i^* small and positive: p^* does not decrease much if we take $v_i > 0$;
if ν_i^* small and negative: p^* does not decrease much if we take $v_i < 0$

local sensitivity: if (in addition) $p^*(u, v)$ is differentiable at $(0, 0)$, then

$$\lambda_i^* = -\frac{\partial p^*(0, 0)}{\partial u_i}, \quad \nu_i^* = -\frac{\partial p^*(0, 0)}{\partial v_i}$$


proof (for λ_i^*): from global sensitivity result,

$$\frac{\partial p^*(0, 0)}{\partial u_i} = \lim_{t \searrow 0} \frac{p^*(te_i, 0) - p^*(0, 0)}{t} \geq -\lambda_i^*$$

$$\frac{\partial p^*(0, 0)}{\partial u_i} = \lim_{t \nearrow 0} \frac{p^*(te_i, 0) - p^*(0, 0)}{t} \leq -\lambda_i^*$$

hence, equality

$p^*(u)$ for a problem with one (inequality) constraint:



Duality and problem reformulations

- equivalent formulations of a problem can lead to very different duals
- reformulating the primal problem can be useful when the dual is difficult to derive, or uninteresting

common reformulations

- introduce new variables and equality constraints
- make explicit constraints implicit or vice-versa
- transform objective or constraint functions

e.g., replace $f_0(x)$ by $\phi(f_0(x))$ with ϕ convex, increasing

Introducing new variables and equality constraints

$$\text{minimize } f_0(Ax + b)$$

- dual function is constant: $g = \inf_x L(x) = \inf_x f_0(Ax + b) = p^*$
- we have strong duality, but dual is quite useless

reformulated problem and its dual

$$\begin{array}{ll} \text{minimize} & f_0(y) \\ \text{subject to} & Ax + b - y = 0 \end{array}$$

$$\begin{array}{ll} \text{maximize} & b^T \nu - f_0^*(\nu) \\ \text{subject to} & A^T \nu = 0 \end{array}$$

dual function follows from

$$\begin{aligned} g(\nu) &= \inf_{x,y} (f_0(y) - \nu^T y + \nu^T Ax + b^T \nu) \\ &= \begin{cases} -f_0^*(\nu) + b^T \nu & A^T \nu = 0 \\ -\infty & \text{otherwise} \end{cases} \end{aligned}$$

norm approximation problem: minimize $\|Ax - b\|$

$$\begin{array}{ll} \text{minimize} & \|y\| \\ \text{subject to} & y = Ax - b \end{array}$$

can look up conjugate of $\|\cdot\|$, or derive dual directly

$$\begin{aligned} g(\nu) &= \inf_{x,y} (\|y\| + \nu^T y - \nu^T Ax + b^T \nu) \\ &= \begin{cases} b^T \nu + \inf_y (\|y\| + \nu^T y) & A^T \nu = 0 \\ -\infty & \text{otherwise} \end{cases} \\ &= \begin{cases} b^T \nu & A^T \nu = 0, \quad \|\nu\|_* \leq 1 \\ -\infty & \text{otherwise} \end{cases} \end{aligned}$$

(see page 5–4)

dual of norm approximation problem

$$\begin{array}{ll} \text{maximize} & b^T \nu \\ \text{subject to} & A^T \nu = 0, \quad \|\nu\|_* \leq 1 \end{array}$$

Implicit constraints

LP with box constraints: primal and dual problem

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & -\mathbf{1} \preceq x \preceq \mathbf{1} \end{array} \qquad \begin{array}{ll} \text{maximize} & -b^T \nu - \mathbf{1}^T \lambda_1 - \mathbf{1}^T \lambda_2 \\ \text{subject to} & c + A^T \nu + \lambda_1 - \lambda_2 = 0 \\ & \lambda_1 \succeq 0, \quad \lambda_2 \succeq 0 \end{array}$$

reformulation with box constraints made implicit

$$\begin{array}{ll} \text{minimize} & f_0(x) = \begin{cases} c^T x & -\mathbf{1} \preceq x \preceq \mathbf{1} \\ \infty & \text{otherwise} \end{cases} \\ \text{subject to} & Ax = b \end{array}$$

dual function

$$\begin{aligned} g(\nu) &= \inf_{-\mathbf{1} \preceq x \preceq \mathbf{1}} (c^T x + \nu^T (Ax - b)) \\ &= -b^T \nu - \|A^T \nu + c\|_1 \end{aligned}$$

dual problem: maximize $-b^T \nu - \|A^T \nu + c\|_1$

Problems with generalized inequalities

$$\begin{array}{ll} \text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \preceq_{K_i} 0, \quad i = 1, \dots, m \\ & h_i(x) = 0, \quad i = 1, \dots, p \end{array}$$

\preceq_{K_i} is generalized inequality on \mathbf{R}^{k_i}

definitions are parallel to scalar case:

- Lagrange multiplier for $f_i(x) \preceq_{K_i} 0$ is vector $\lambda_i \in \mathbf{R}^{k_i}$
- Lagrangian $L : \mathbf{R}^n \times \mathbf{R}^{k_1} \times \dots \times \mathbf{R}^{k_m} \times \mathbf{R}^p \rightarrow \mathbf{R}$, is defined as

$$L(x, \lambda_1, \dots, \lambda_m, \nu) = f_0(x) + \sum_{i=1}^m \lambda_i^T f_i(x) + \sum_{i=1}^p \nu_i h_i(x)$$

- dual function $g : \mathbf{R}^{k_1} \times \dots \times \mathbf{R}^{k_m} \times \mathbf{R}^p \rightarrow \mathbf{R}$, is defined as

$$g(\lambda_1, \dots, \lambda_m, \nu) = \inf_{x \in \mathcal{D}} L(x, \lambda_1, \dots, \lambda_m, \nu)$$

lower bound property: if $\lambda_i \succeq_{K_i^*} 0$, then $g(\lambda_1, \dots, \lambda_m, \nu) \leq p^*$

proof: if \tilde{x} is feasible and $\lambda \succeq_{K_i^*} 0$, then

$$\begin{aligned} f_0(\tilde{x}) &\geq f_0(\tilde{x}) + \sum_{i=1}^m \lambda_i^T f_i(\tilde{x}) + \sum_{i=1}^p \nu_i h_i(\tilde{x}) \\ &\geq \inf_{x \in \mathcal{D}} L(x, \lambda_1, \dots, \lambda_m, \nu) \\ &= g(\lambda_1, \dots, \lambda_m, \nu) \end{aligned}$$

minimizing over all feasible \tilde{x} gives $p^* \geq g(\lambda_1, \dots, \lambda_m, \nu)$

dual problem

$$\begin{aligned} &\text{maximize} && g(\lambda_1, \dots, \lambda_m, \nu) \\ &\text{subject to} && \lambda_i \succeq_{K_i^*} 0, \quad i = 1, \dots, m \end{aligned}$$

- weak duality: $p^* \geq d^*$ always
- strong duality: $p^* = d^*$ for convex problem with constraint qualification (for example, Slater's: primal problem is strictly feasible)

Semidefinite program

primal SDP ($F_i, G \in \mathbf{S}^k$)

$$\begin{aligned} & \text{minimize} && c^T x \\ & \text{subject to} && x_1 F_1 + \cdots + x_n F_n \preceq G \end{aligned}$$

- Lagrange multiplier is matrix $Z \in \mathbf{S}^k$
- Lagrangian $L(x, Z) = c^T x + \mathbf{tr}(Z(x_1 F_1 + \cdots + x_n F_n - G))$
- dual function

$$g(Z) = \inf_x L(x, Z) = \begin{cases} -\mathbf{tr}(GZ) & \mathbf{tr}(F_i Z) + c_i = 0, \quad i = 1, \dots, n \\ -\infty & \text{otherwise} \end{cases}$$

dual SDP

$$\begin{aligned} & \text{maximize} && -\mathbf{tr}(GZ) \\ & \text{subject to} && Z \succeq 0, \quad \mathbf{tr}(F_i Z) + c_i = 0, \quad i = 1, \dots, n \end{aligned}$$

$p^* = d^*$ if primal SDP is strictly feasible ($\exists x$ with $x_1 F_1 + \cdots + x_n F_n \prec G$)