

8. Geometric problems

- extremal volume ellipsoids
- centering
- classification
- placement and facility location

Minimum volume ellipsoid around a set

Löwner-John ellipsoid of a set C : minimum volume ellipsoid \mathcal{E} such that $C \subseteq \mathcal{E}$

- parametrize \mathcal{E} as $\mathcal{E} = \{v \mid \|Av + b\|_2 \leq 1\}$; w.l.o.g. assume $A \in \mathbf{S}_{++}^n$
- **vol** \mathcal{E} is proportional to $\det A^{-1}$; to compute minimum volume ellipsoid,

$$\begin{array}{ll} \text{minimize (over } A, b) & \log \det A^{-1} \\ \text{subject to} & \sup_{v \in C} \|Av + b\|_2 \leq 1 \end{array}$$

convex, but evaluating the constraint can be hard (for general C)

Finite set $C = \{x_1, \dots, x_m\}$:

$$\begin{array}{ll} \text{minimize (over } A, b) & \log \det A^{-1} \\ \text{subject to} & \|Ax_i + b\|_2 \leq 1, \quad i = 1, \dots, m \end{array}$$

also gives Löwner-John ellipsoid for polyhedron $\text{conv}\{x_1, \dots, x_m\}$

Maximum volume inscribed ellipsoid

maximum volume ellipsoid \mathcal{E} inside a convex set $C \subseteq \mathbf{R}^n$

- parametrize \mathcal{E} as $\mathcal{E} = \{Bu + d \mid \|u\|_2 \leq 1\}$; w.l.o.g. assume $B \in \mathbf{S}_{++}^n$
- $\text{vol } \mathcal{E}$ is proportional to $\det B$; can compute \mathcal{E} by solving

$$\begin{aligned} & \text{maximize} && \log \det B \\ & \text{subject to} && \sup_{\|u\|_2 \leq 1} I_C(Bu + d) \leq 0 \end{aligned}$$

(where $I_C(x) = 0$ for $x \in C$ and $I_C(x) = \infty$ for $x \notin C$)

convex, but evaluating the constraint can be hard (for general C)

Polyhedron $\{x \mid a_i^T x \leq b_i, i = 1, \dots, m\}$:

$$\begin{aligned} & \text{maximize} && \log \det B \\ & \text{subject to} && \|Ba_i\|_2 + a_i^T d \leq b_i, \quad i = 1, \dots, m \end{aligned}$$



(constraint follows from $\sup_{\|u\|_2 \leq 1} a_i^T (Bu + d) = \|Ba_i\|_2 + a_i^T d$)

Efficiency of ellipsoidal approximations

$C \subseteq \mathbf{R}^n$ convex, bounded, with nonempty interior

- Löwner-John ellipsoid, shrunk by a factor n , lies inside C
- maximum volume inscribed ellipsoid, expanded by a factor n , covers C

Example (for two polyhedra in \mathbf{R}^2)





factor n can be improved to \sqrt{n} if C is symmetric

Centering

some possible definitions of 'center' of a convex set C :

- center of largest inscribed ball ('Chebyshev center')
for polyhedron, can be computed via linear programming (page 4.16)
- center of maximum volume inscribed ellipsoid (page 8.3)



MVE center is invariant under affine coordinate transformations

Analytic center of a set of inequalities

the analytic center of set of convex inequalities and linear equations

$$f_i(x) \leq 0, \quad i = 1, \dots, m, \quad Fx = g$$

is defined as the optimal point of

$$\begin{array}{ll} \text{minimize} & - \sum_{i=1}^m \log(-f_i(x)) \\ \text{subject to} & Fx = g \end{array}$$


- more easily computed than MVE or Chebyshev center (see later)
- not just a property of the feasible set: two sets of inequalities can describe the same set, but have different analytic centers

Analytic center of linear inequalities

$$a_i^T x \leq b_i, \quad i = 1, \dots, m$$

x_{ac} is minimizer of

$$\phi(x) = - \sum_{i=1}^m \log(b_i - a_i^T x)$$



inner and outer ellipsoids from analytic center:

$$\mathcal{E}_{\text{inner}} \subseteq \{x \mid a_i^T x \leq b_i, i = 1, \dots, m\} \subseteq \mathcal{E}_{\text{outer}}$$

where


$$\mathcal{E}_{\text{inner}} = \{x \mid (x - x_{ac})^T \nabla^2 \phi(x_{ac})(x - x_{ac}) \leq 1\}$$

$$\mathcal{E}_{\text{outer}} = \{x \mid (x - x_{ac})^T \nabla^2 \phi(x_{ac})(x - x_{ac}) \leq m(m - 1)\}$$

Linear discrimination

separate two sets of points $\{x_1, \dots, x_N\}$, $\{y_1, \dots, y_M\}$ by a hyperplane:

$$a^T x_i + b > 0, \quad i = 1, \dots, N, \quad a^T y_i + b < 0, \quad i = 1, \dots, M$$



homogeneous in a , b , hence equivalent to

$$a^T x_i + b \geq 1, \quad i = 1, \dots, N, \quad a^T y_i + b \leq -1, \quad i = 1, \dots, M$$

a set of linear inequalities in a , b


Robust linear discrimination

(Euclidean) distance between hyperplanes

$$\mathcal{H}_1 = \{z \mid a^T z + b = 1\}$$

$$\mathcal{H}_2 = \{z \mid a^T z + b = -1\}$$

is $d(\mathcal{H}_1, \mathcal{H}_2) = 2/\|a\|_2$



to separate two sets of points by maximum margin,

$$\begin{aligned} & \text{minimize} && (1/2)\|a\|_2 \\ & \text{subject to} && a^T x_i + b \geq 1, \quad i = 1, \dots, N \\ & && a^T y_i + b \leq -1, \quad i = 1, \dots, M \end{aligned} \tag{1}$$

(after squaring objective) a QP in a, b

Lagrange dual of maximum margin separation problem (1)

$$\begin{aligned} & \text{maximize} && \mathbf{1}^T \lambda + \mathbf{1}^T \mu \\ & \text{subject to} && 2 \left\| \sum_{i=1}^N \lambda_i x_i - \sum_{i=1}^M \mu_i y_i \right\|_2 \leq 1 \\ & && \mathbf{1}^T \lambda = \mathbf{1}^T \mu, \quad \lambda \geq 0, \quad \mu \geq 0 \end{aligned} \tag{2}$$

from duality, optimal value is inverse of maximum margin of separation

Interpretation

- change variables to

$$\theta_i = \frac{\lambda_i}{\mathbf{1}^T \lambda}, \quad \gamma_i = \frac{\mu_i}{\mathbf{1}^T \mu}, \quad t = \frac{1}{\mathbf{1}^T \lambda + \mathbf{1}^T \mu}$$

- invert objective to minimize $1/(\mathbf{1}^T \lambda + \mathbf{1}^T \mu) = t$


$$\begin{aligned} & \text{minimize} && t \\ & \text{subject to} && \left\| \sum_{i=1}^N \theta_i x_i - \sum_{i=1}^M \gamma_i y_i \right\|_2 \leq t \\ & && \theta \geq 0, \quad \mathbf{1}^T \theta = 1, \quad \gamma \geq 0, \quad \mathbf{1}^T \gamma = 1 \end{aligned}$$

optimal value is distance between convex hulls

Approximate linear separation of non-separable sets

$$\begin{aligned} & \text{minimize} && \mathbf{1}^T u + \mathbf{1}^T v \\ & \text{subject to} && a^T x_i + b \geq 1 - u_i, \quad i = 1, \dots, N \\ & && a^T y_i + b \leq -1 + v_i, \quad i = 1, \dots, M \\ & && u \geq 0, \quad v \geq 0 \end{aligned}$$

- an LP in a, b, u, v
- at optimum, $u_i = \max\{0, 1 - a^T x_i - b\}$, $v_i = \max\{0, 1 + a^T y_i + b\}$
- can be interpreted as a heuristic for minimizing #misclassified points




Support vector classifier

$$\begin{aligned} & \text{minimize} && \|a\|_2 + \gamma(\mathbf{1}^T u + \mathbf{1}^T v) \\ & \text{subject to} && a^T x_i + b \geq 1 - u_i, \quad i = 1, \dots, N \\ & && a^T y_i + b \leq -1 + v_i, \quad i = 1, \dots, M \\ & && u \geq 0, \quad v \geq 0 \end{aligned}$$

produces point on trade-off curve between inverse of margin $2/\|a\|_2$ and classification error, measured by total slack $\mathbf{1}^T u + \mathbf{1}^T v$

same example as previous page,
with $\gamma = 0.1$:



Nonlinear discrimination

separate two sets of points by a nonlinear function:

$$f(x_i) > 0, \quad i = 1, \dots, N, \quad f(y_i) < 0, \quad i = 1, \dots, M$$

- choose a linearly parametrized family of functions

$$f(z) = \theta^T F(z)$$

$F = (F_1, \dots, F_k) : \mathbf{R}^n \rightarrow \mathbf{R}^k$ are basis functions

- solve a set of linear inequalities in θ :


$$\theta^T F(x_i) \geq 1, \quad i = 1, \dots, N, \quad \theta^T F(y_i) \leq -1, \quad i = 1, \dots, M$$

Quadratic discrimination: $f(z) = z^T P z + q^T z + r$


$$x_i^T P x_i + q^T x_i + r \geq 1, \quad y_i^T P y_i + q^T y_i + r \leq -1$$

can add additional constraints (e.g., $P \leq -I$ to separate by an ellipsoid)

Polynomial discrimination: $F(z)$ are all monomials up to a given degree



separation by ellipsoid



separation by 4th degree polynomial

Placement and facility location

- N points with coordinates $x_i \in \mathbf{R}^2$ (or \mathbf{R}^3)
- some positions x_i are given; the other x_i 's are variables
- for each pair of points, a cost function $f_{ij}(x_i, x_j)$

Placement problem

$$\text{minimize } \sum_{i \neq j} f_{ij}(x_i, x_j)$$




variables are positions of free points

Interpretations

- points represent plants or warehouses; f_{ij} is transportation cost between facilities i and j
- points represent cells on an IC; f_{ij} represents wirelength

Example: minimize $\sum_{(i,j) \in \mathcal{A}} h(\|x_i - x_j\|_2)$, with 6 free points, 27 links

optimal placement for $h(z) = z$, $h(z) = z^2$, $h(z) = z^4$



histograms of connection lengths $\|x_i - x_j\|_2$

